

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pathology and Oncology

Journal homepage: [www.ijpo.co.in](http://www.ijpo.co.in)

## Original Research Article

## Hypersegmented neutrophils in peripheral smear –An etiological analysis

Feby T Francis<sup>1,\*</sup>, Aysha Ali<sup>1</sup>, Priya P Kartha<sup>1</sup>, Jayasree K A<sup>1</sup><sup>1</sup>Dept. of Pathology, Government Medical College, Palakkad, Kerala, India

## ARTICLE INFO

## Article history:

Received 08-12-2021

Accepted 16-12-2021

Available online 14-02-2022

## Keywords:

Microcytic hypochromic anemia

Neutrophil hypersegmentation

Thrombocytosis

## ABSTRACT

Hypersegmentation of neutrophils is defined as presence of 5% or more neutrophils with five or more lobes or single neutrophil with 6 lobes. It is usually associated with deficiency of or failure to utilize cobalamin or folate and impaired DNA synthesis is the accepted mechanism for the morphological changes seen in megaloblastosis. Other causes of neutrophil hypersegmentation (NH) listed include microcytic hypochromic anemia (MHA) but the evidence for this is based mainly on a limited number of case studies in which MHAs with vit B12 and folic acid deficiency also were included. More studies are needed to establish etiological factors other than already established megaloblastic anemia.

The present study was conducted in Department of Pathology in a tertiary care centre in Central Kerala, India. 100 cases peripheral smears with NH were evaluated. Aim of the study was to classify the etiological factors of NH in peripheral smears, study NH in microcytic hypochromic anemia and to check whether there is any connection between microcytic hypochromic anemia with NH and thrombocytosis. Complete blood count of all cases were taken using automated hematology analysers. Peripheral smear picture were correlated with the blood counts. Cases with MHA were examined for underlying vit B12 and folic acid deficiency. Study shows that 31% cases of peripheral smears with hypersegmentation of neutrophils were pure microcytic hypochromic anemias with normal vitB12 and folic acid levels. Study also points to increased incidence thrombocytosis in pure MHA cases compared to other etiological factors.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Hypersegmentation of neutrophils is defined as presence of 5% or more neutrophils with five or more lobes or single neutrophil with 6 lobes.<sup>1</sup> It is usually associated with deficiency of or failure to utilize cobalamin or folate and impaired DNA synthesis is the accepted mechanism for the morphological changes seen in megaloblastosis.<sup>2,3</sup> Other causes of NH listed include microcytic hypochromic anemia but the evidence for this is based mainly on a limited number of case studies in which patients in whom the IDA complicated by coexistent cobalamin or folate deficiency were also included.<sup>4,5</sup> Hypersegmented neutrophils are also

seen as a part of myelodysplastic syndromes which is usually designated as bone marrow (BM) failure are a heterogeneous group of myeloid clonal disorders caused by a failure of blood cells maturation. The co-morbidities result from a variable degree of cytopenia and clonal instability with a tendency to progression mainly into acute myeloid leukemia (AML).<sup>6</sup> Uremia, hyperthermia, drugs including chemotherapy, steroids, GCSF are also known to produce neutrophil hypersegmentation.<sup>7</sup> It is also known in a congenital condition (autosomal dominant) affecting 1% of the population.<sup>8</sup>

Here in the present study we evaluated 100 peripheral smears with hypersegmented neutrophils and classified the etiological factors. Patients with microcytic hypochromic anemia were further evaluated for underlying vit B12 and

\* Corresponding author.

E-mail address: [drchristytf@gmail.com](mailto:drchristytf@gmail.com) (F. T. Francis).

folic acid deficiency. This study also checks whether there is any association between neutrophil hypersegmentation in microcytic hypochromic anemia and thrombocytosis.

## 2. Material and Methods

This is a prospective study starting from 2020 January to 2020 December conducted in Department of Hematology in a tertiary care centre in central Kerala. EDTA blood samples received in our hematology laboratory were analysed for hypersegmentation of neutrophils using geimsa stained peripheral smears. Neutrophils hypersegmentation is defined as the presence of five or more five-lobed neutrophils per 100, or any neutrophils with six or more lobes. 100 such cases which satisfied the inclusion criteria were taken as sample size. Complete blood count of individual cases were obtained using Sysmex SE9000 analyser and peripheral smear picture was compared with blood counts. These 100 cases were classified according to the peripheral smear picture. Patients with microcytic hypochromic anemia were separately assessed for serum Vit B12 and folic acid values using ion capture assay and microparticle enzyme intrinsic factor assay. Presence of thrombocytosis in pure microcytic hypochromic anemia cases were checked separately and it was compared with presence of thrombocytosis in cases with NH without microcytic hypochromic anemia. Patients with known medical conditions like pregnancy, uremia, renal failure and exposure to drugs like chemotherapy, steroid and GCSF were excluded.

## 3. Results

Detailed analysis of all cases showed the following details. Age and gender distribution of cases are shown in Table 1. Majority of cases were males and majority of cases were in the age group 40-60.

Cases were further analysed for associated peripheral smear picture. Although major cases were contributed by macrocytic anemia, 41% cases were having microcytic hypochromic anemia. Detailed picture is given in Table 2.

Detailed analysis of peripheral smears were done to check whether neutrophil hypersegmentation was associated with any other neutrophil abnormalities. 22 cases showed cytoplasmic toxic granules along with nuclear hypersegmentation. 18 cases showed cytoplasmic vacuoles. 5 cases showed both toxic granules and vacuoles. 8 cases showed Dohle bodies. One case among 3 myelodysplastic syndrome case showed hypogranular neutrophil. Rest of the 46 cases didn't show any neutrophil abnormality other than hypersegmentation.

Changes in the counts of white blood cells were also evaluated along with neutrophil hypersegmentation. 34% of cases showed neutrophilia. Eosinophilia was seen in 8% of cases. No cases showed lymphocytosis, monocytosis or basophilia.

41 cases of neutrophil hypersegmentation which showed microcytic hypochromic blood picture were checked for underlying vit B12 and folic acid deficiency. Normal range of Serum vit B12 and folic acid values are 200-900pg/ml and 2-20ng/ml.

Table 3 clearly shows that out of the 41 cases with normocytic normochromic bloodpicture, only 10 had subnormal levels of either Vit B12 or folic acid values. Rest of the 30 cases had normal Vit B12 and folic acid levels. So we can come to the conclusion that out of the 100 cases with hypersegmented neutrophils in peripheral smear 31% cases were having pure microcytic hypochromic anemia without any vit B12 or folic acid deficiency.

Platelet count of all cases were assessed. Results are shown in Table 4.

1.5-4.5 lakh/microliter is considered as normal platelet count. Out of the 100 cases, only 2 had thrombocytopenia. 73 cases had platelet count in the normal range. 25 cases had thrombocytosis. Out of the 31 cases with microcytic hypochromic anemia and neutrophil hypersegmentation, 19 cases had thrombocytosis (61.29%). In all other cases majority were in normal range group.

## 4. Discussion

In this study of 100 cases with peripheral smears, showed that there are many causes for neutrophil hypersegmentation other than already established macrocytic anemia. Microcytic hypochromic anemia, melodysplastic syndrome and normocytic normochromic blood picture show neutrophil hypersegmentation in peripheral smear. Deficiency of or failure to utilize cobalamin or folate and impaired DNA synthesis is the accepted mechanism for the morphological changes seen in megaloblastosis.<sup>2,3</sup>

The underlying mechanism of neutrophil hypersegmentation in microcytic hypochromic anemia is not fully understood. There are several studies explaining this as undetected Vit B12 and folic acid deficiency.<sup>4,5</sup> But that is unlikely as 31% cases of NH were pure microcytic hypochromic anemia without Vit B12 and folic acid deficiency. As per previous studies which excluded underlying Vit B12 and folic acid deficiency there are significant association between iron deficiency (presenting as microcytic hypochromic anemia) and neutrophil hypersegmentation.<sup>3</sup> There are some explanations in previous studies that neutrophil changes in iron deficiency represent a recent event with only young red cells and not the overall population of red cells, showing a reduced folate content.<sup>9,10</sup> Some other studies shows that iron deficiency can affect the folate dependant degradation of Figlu catalysed by enzyme Figlu transferase.<sup>11,12</sup> It is also possible that iron deficiency may directly influence DNA synthesis.<sup>13,14</sup>

**Table 1:** Age and gender distribution of all cases showing hyper segmented neutrophils in peripheral smears

Gender	Below 20	20-40	40-60	Above 60	Total
Male	9	11	19	15	54
Female	5	9	20	12	46
Total	14	20	39	27	100

**Table 2:** Peripheral smear picture of cases with hyper segmented neutrophils

Macrocytic anemia	45
Microcytic hypochromic anemia	41
Normocytic normochromic blood picture	11
Myelodysplastic syndrome	3
Total	100

**Table 3:** Serum Vit B12 and folic acid values of cases with neutrophil hypersegmentation in microcytic hypochromic blood picture

Vit B12(in pg/ml)	Observed frequency	Folic acid (in ng/ml)	Observed frequency
<200pg/ml	7	<2ng/ml	3
200-500pg/ml	22	2-8ng/ml	8
500-700pg/ml	9	8-15ng/ml	19
700-900pg/ml	2	15-20ng/ml	11
>900pg/ml	1	>20ng/ml	0
Total	41		41

**Table 4:** Correlation of neutrophil hypersegmentation and platelet count

Platelet count	Macrocytic anemia	Microcytic hypochromic Picture (Normal B12 and folic acid)	Microcytic hypochromic Picture (subnormal B12 and folic acid)	Myelodysplasia	Normocytic Normochromic Blood picture
<1.5 lakh/microlitre	1	0	0	1	0
1.5-4.5 lakh/microlitre	41	12	9	2	9
>4.5lakh/microlitre	3	19	1	0	2
Total	45	31	10	3	11

There are several studies showing that hypersegmented neutrophils can be seen as a part of trauma and chronic infections. It is also said to be recruited to bloodstream during inflammation.<sup>15</sup> In our study 22 cases showed neutrophil toxic granules along with hypersegmentation. 18 cases showed vacuoles also. Both toxic granules and vacuoles are known to be the response to infection, inflammation and stress.<sup>16</sup> The presence of toxic granules and vacuoles along with hypersegmentation in our study also point towards the appearance of hypersegmentation as a part of inflammation.

3 cases in our study were diagnosed as myelodysplastic syndromes. There are several earlier studies demonstrating that neutrophil hypersegmentation can be seen as a part of myelodysplasia.<sup>17,18</sup> One case of myelodysplastic syndrome in this study showed hypolobated neutrophil along with hyper segmented neutrophils.

Neutrophil hypersegmentation and thrombocytosis can be expected as a part of megaloblastic anemia. But it may be also seen in iron deficiency anemia also.<sup>19,20</sup> In the present

study out of the 25 cases having thrombocytosis (>4.5lakhs/ml), 19 cases are having microcytic hypochromic anemia with normal vit B12 and folic acid values. But further detailed studies are to be done to know whether there is any association between neutrophil hypersegmentation and thrombocytosis.

## 5. Conclusion

The present study indicates that other than the already established causes of neutrophil hypersegmentation, microcytic hypochromic anemia, myelodysplastic syndromes and inflammatory conditions also can cause hypersegmented neutrophils in peripheral smears. Study also points to increased incidence thrombocytosis in pure MHA cases compared to other etiological factors which is still to be established with further detailed studies.

## 6. Source of Funding

None.

## 7. Conflict of Interest

The authors declare no conflict of interest.

## References

- Herbert V. Nutrition science as a continually unfolding story: the folate and vitamin B12 paradigm. *Am J Clin Nutr.* 1987;46(3):387–402.
- Agarwal KN. Indicators for assessment of anaemia and iron deficiency in community. *Pediatr Oncall J.* 2010;7(2):29–34.
- Westerman DDA, Evans D, Metz J. Neutrophil hypersegmentation in iron deficiency anaemia: A case control study. *Br J Haematol.* 1999;107(3):512–5.
- Doscherholmen KA, Muhamud D, Ripley. Hypersegmentation in iron deficiency anaemia. *JAMA.* 1974;229(13):1721–2.
- Rao GB, Lokeswari V, Prasad PR. Hypersegmentation of neutrophils—still a hallmark of macrocytic anemia? A prospective study. *Int J Sci Res.* 2018;7(4):49–50.
- Steensma DP. Myelodysplastic syndromes: diagnosis and treatment. *Mayo Clin Proc.* 2015;90(7):969–83.
- Im DD, Ho CH, Chan RY. Hypersegmented Neutrophils in an Adolescent Male With Heatstroke. *J Pediatr Hematol Oncol.* 2015;37(6):488.
- Shinton NK. Desk Reference for Hematology. 2nd ed. United States: CRC Press; 2007.
- Omer A, Finlayson ND, Shearman DJC, Samson RR, Girdwood RH. Plasma and erythrocyte folate in iron deficiency and folate deficiency. *Blood.* 1970;35(6):821–8.
- Roberts PD, John DS, Sinha R, Stewart JS, Baird IM, Coghill NF, et al. Apparent folate deficiency in iron deficiency anemia. *Br J Haematol.* 1971;20(2):165–75.
- Chanarin I, Bennett MC, Berry V. Urinary excretion of histidine derivative in megaloblastic anemia and other conditions in comparison with the folic acid clearance test. *J Clin Pathol.* 1962;15(3):269–73.
- Vitale JJ, Restrepo A, Riker JB, Hellerstin EE. Secondary folate deficiency induced in rat by dietary iron deficiency. *J Nutr.* 1966;88(3):315–22.
- Hershko CH, Karsai A, Eylon L, Izak G. The effect of chronic iron deficiency on some biochemical functions of the human hematopoietic tissue. *Blood.* 1970;36(3):321–9.
- Pillay J, Kamp VM, Hoffen EV, Visser T, Tak T, Lammers J. A subset of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1. *J Clin Invest.* 2012;122:327–36.
- Tak T, Wijten P, Heeres M, Pickkers P, Scholten A, Heck A. Human CD62L dim neutrophils identified as a separate subset by proteome profiling and in vivo pulse-chase labelling. *Blood.* 2017;129:3476–86.
- Manonneaux S. Non malignant leukocyte disorders. In: Rodaks hematology clinical applications and principals. St. Louis, Missouri: Saunders; 2015. p. 475–97.
- Yoshida Y. Physical Education. Myelodysplastic syndrome. *Oncologist.* 1996;1(4):284–7.
- Abramson SD, Abramson N. Common 'uncommon' anemias. *Am Fam Physician.* 1999;59(4):851–8.
- Duzgun S, Yildirmak Y, Cetinkaya F. Neutrophil hypersegmentation and thrombocytosis in children with iron deficiency anemia. *Turk J Pediatr.* 2005;47:251–4.
- Mahmood T, Robinson WA, Hamstra RD, Wallner SF. Macrocytic anemia, thrombocytosis and non lobulated megakaryocytes: the 5q-syndrome, a distinct entity. *Am J Med.* 1979;66(6):946–50.

## Author biography

**Feby T Francis**, Assistant Professor

**Aysha Ali**, Associate Professor

**Priya P Kartha**, Assistant Professor

**Jayasree K A**, Assistant Professor

**Cite this article:** Francis FT, Ali A, Kartha PP, Jayasree K A. Hypersegmented neutrophils in peripheral smear –An etiological analysis. *Indian J Pathol Oncol* 2022;9(1):39-42.