



Original Research Article

The role of hematological biomarkers, coagulation profile and CRP (c-reactive protein) in COVID-19 patients in private health care sector

Sneha Aditi^{1,*}, Roushni Kumari², Abdul Aziz¹¹SRL Diagnosis, Patna, Bihar, India²Nalanda Medical College and Hospital, Patna, Bihar, India

ARTICLE INFO

Article history:

Received 29-06-2022

Accepted 19-07-2022

Available online 26-08-2022

Keywords:

SARS- CoV- 2

RT-PCR

Systemic inflammatory index

Neutrophil lymphocyte ratio

Platelet lymphocyte ratio

Coagulation factor

Coronavirus

ABSTRACT

Background: SARS-CoV-2 causing corona virus disease 2019 (COVID-19) is the most common infection in recent days worldwide. Rapidly evolving from the epidemic outbreak in Wuhan, China on 29 December 2019, SARS-CoV-2 is the third known corona virus that causes fatal respiratory diseases in humans.

Aim: The roles of biomarkers from a blood sample in the diagnosis of COVID-19 patients who have visited the hospital with history of fever has been evaluated.

Materials and Methods: A retrospective study was carried out in a private sector hospital at Panipat where the patients having history of fever were admitted during 1st May 2020 to 15th August 2020. A total of 100 patients were included who were suspected cases of corona. All the blood parameters, systemic inflammatory index (SII), neutrophil lymphocyte ratio (NLR) and platelet lymphocyte ratio (PLR) Coagulation factor (prothrombin time (PT), activated partial thromboplastin time (aPTT), international normalized ratio (INR) and C-Reactive protein (CRP) are compared between negative and positive COVID-19 cases according to RT-PCR. Result: Out of 100 patients, 65% were RTPCR covid positive. In patient with RTPCR negative test results, it was found that total leukocyte count, neutrophils, platelet, NLR and SII values were higher. However, haemoglobin, lymphocytes and PLR was found to be elevated in patients with positive test results.

Discussion: COVID-19 positive patients presented with leukopenia, neutropenia, thrombocytopenia and lymphocytosis. SII and NLR were reduced. PLR and CRP were increased. PT and APTT were within normal limits. Lymphocyte count and PLR showed strong significance in this study.

Conclusion: The definitive diagnosis of COVID-19 was made by RT-PCR analysis, but this is a time-consuming and less accessible test. Low values of NLR and SII and high value of PLR and CRP are indicative of COVID-19 and can be used in its diagnosis.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

A bunch of mysterious pneumonia cases was reported by the Chinese population to the World Health Organization (WHO) on 31 December, 2019.¹ The etiology for this outbreak was a novel corona virus named severe acute respiratory syndrome corona virus 2 (SARS-CoV-2), which was responsible for the Corona Virus Disease 2019

(COVID-19).² And by January 2020 the disease became epidemic and was observed internationally, cases have been reported in 24 countries and 5 continents.³

Although it is well expected that COVID-19 is primarily manifested as a respiratory tract infection, up-and-coming data indicate that it should be regarded as a systemic disease involving multiple systems, including cardiovascular, respiratory, gastrointestinal, neurological and hematopoietic and immune system.⁴ Patient's clinical manifestations

* Corresponding author.

E-mail address: 25.adisneha@gmail.com (S. Aditi).

included fever, nonproductive cough, dyspnea, myalgia, fatigue and radiological substantiation of pneumonia. Organ dysfunction (e.g., shock, acute respiratory distress syndrome [ARDS], acute cardiac injury, and acute kidney injury) and death can occur in severe cases.¹ Early diagnosis is vital when bearing in mind the short time of onset of acute respiratory distress syndrome among patients after admission to hospital and the high mortality rates in COVID-19.

Blood tests have an important role in early diagnosis of the disease, considering the information they provide to physicians regarding the inflammatory process. Complete blood counts (CBC) are easily performed and cheap. Included in the CBC are values such as white blood count, neutrophil, lymphocyte and platelet count (PLTs). These parameters may be used as inflammatory markers by themselves, their ratios to one another may also be indicators of early inflammation.¹ Considering previous research, the use of circulating biomarkers in lieu of inflammation and the immune system have been considered as a prognostic indicator in COVID-19-positive patients.

In this study, the roles of biomarkers from a peripheral blood sample is used in the diagnosis of COVID-19 patients who have history of fever and have been admitted to hospital are examined.

2. Materials and Methods

A retrospective study was carried out in a private sector hospital at Panipat where the patients having history of fever were admitted during 1st May 2020 to 15th August 2020. A total of 100 patients were included who were suspected cases of corona. RTPCR was done for all the patients and were categorized into covid negative and covid positive patients.


All the blood parameters, Systemic inflammatory index (SII) = thrombocyte count \times neutrophil count/lymphocyte count. Neutrophil lymphocyte ratio (NLR) = absolute neutrophil count/absolute lymphocyte count. Platelet lymphocyte ratio = absolute platelet/absolute lymphocyte count were calculated. Coagulation factor (prothrombin time(PT), activated partial thromboplastin time (aPTT), international normalized ratio(INR) and C-Reactive protein (CRP) were compared between negative and positive COVID-19 cases.

The analysis of the data was done using the SPSS 25 statistical package programs. A Chi-square test was used to analyze the resounding variables of the patients, which were expressed as a number and percentage. The independent samples t-test was used for analysis for parametric continued variables and they were presented as a mean as well as standard deviation. The Binary Logistic Regression test was used to assess the diagnostic utility of several biomarkers to define risk ratios, including the SII. A receiver-operating characteristic (ROC) curve was

formed, the Youden's index (J) used for cut-off values in the diagnosis of COVID-19 and the area under the curve (AUC) was calculated, $p < 0.05$ was considered statistically significant.¹

3. Results

Out of 100 patients included in the study that came with history of fever and was admitted in hospital, RTPCR was done and 65% patients were positive. Out of which 66.2% were male patients and 33.8% were female patients as shown in Graph 1. The mean age of positive patients was 43.03years with standard deviation of 16.1.



Graph 1: Covid suspected patients vs sex

Hematological parameters were recorded and presented in a tabular data as described in Table 1. Parameters included were total leukocyte count, neutrophils, hemoglobin, lymphocytes, NLR, PLR, SII, and Platelets. In patient with negative test results, it was found that total leukocyte count, neutrophils, platelet, NLR and SII values were higher. Whereas haemoglobin, lymphocytes and PLR was found to be higher in patients with positive test results.

Since there was a statistically significant disparity between negative and positive groups according to parameters (leukocyte, neutrophil hemoglobin, platelet, NLR, PLR and SII), they were then studied with ROC analyses. Regarding the observations made by ROC analyses, the following information concerning patients with COVID-19 diagnoses were found (Table 1 & Figure 1). As shown in Figure 1 hemoglobin, lymphocyte and PLR was noted towards the positive side of the curve.

TLC, Neutrophils, platelets, SII and NLR could not be used as a potential diagnostic biomarker for subsequent analysis because its AUC was less than 0.50. The optimal cut-off values were 11.25, 22.5 and 201.8 for hemoglobin, lymphocytes and PLR.

The optimal cut-off values were 11.25, 22.5 and 201.8 for hemoglobin, lymphocytes and PLR. Comparisons were made between these two groups that were formed according to the cut-off values and correlation was established with PLR where p value was < 0.05 .

Further coagulation profile (PT, APTT, INR) and CRP was compared between negative and positive group. P value was assessed and there was strong correlation noted with

Table 1: Comparison of haematological parameters according to RTPCR test results

Hematological parameters	Positive (n=65)	Negative (n=35)
Total leukocyte count	9.7±5.7 7.7(3.8-26.1)	12.5±5.3 13.1(5.3-23.2)
Neutrophils	6.2± 1.3 6.2(3.2-9.3)	6.8±1.3 6.7(3.5-9)
Hemoglobin	13.3±1.5 13.4 (9.7-16.5)	11.8±2.6 12.4(6-16)
Lymphocytes	2.6±1.2 2.7(0.4-5.8)	2.2±1.1 2(0.5-5.3)
NLR	3.6±3.6 2.3(0.55-23.25)	4.6±3.9 3.2(0.66-18)
PLR	130.27±85.87 112.17(20.4-383.26)	119.89±63.7 104.6(29.5-262.3)
SII	861.4±120 457(111-862.5)	999.6±78.67 748(311-4014)
Platelet	224.2 ± 92.7 212.0(79- 458)	241±97.8 212(95-510)

Table 2: Area under the curve for hemoglobin, TLC, Neutrophils, lymphocytes, platelets, SII, NLR and PLR

Test Result Variable(s)	Area	Std. Error ^a	Asymptotic Sig. ^b	Asymptotic 95% Confidence Interval	
				Lower Bound	Upper Bound
Hemoglobin	.671	.058	.005	.557	.785
TLC	.332	.054	.006	.225	.438
Neutrophils	.367	.057	.029	.256	.478
Lymphocytes	.610	.057	.070	.498	.722
Platelets	.447	.060	.380	.329	.564
SII	.346	.054	.011	.241	.451
NLR	.382	.057	.053	.271	.494
PLR	.506	.059	.917	.390	.623

Table 3: Comparison of results according to cut-off points

Hematological parameters		Negative	Positive	p value
Hemoglobin	<11.25	14(40%)	6(9.2%)	0.06
	>11.25	21(60%)	59(90.76%)	
lymphocytes	<22.5	18(51.4%)	20(30.7%)	0.05
	>22.5	17(48.57%)	45(69.2%)	
PLR	<201.8	29(82.85%)	52(80%)	0.03
	>201.8	6(17.14)	13(20%)	

Table 4: Comparison of coagulation profile and CRP

Coagulation profile and CRP	Positive	Negative	P value
PT (Prothrombin time)	14.33±.44	14.54±0.644	.187
APTT (Activated partial thromboplastin time)	30±0.01	31.50±1.3	.117
INR	1.3±0.07	1.1±0.05	.06
CRP	7.9±0.26	6.03±0.5	.05

CRP as shown in Table 4.

4. Discussion


Taking into account that COVID 19 is highly infective and has high mortality rates early diagnosis of the disease is essential. The authoritative diagnosis of this disease is made by a viral presence in real-time PCR analyses. Due to factors such as the high number of samples, limited number of staff trained in performing the aforementioned tests insufficient lab capacities, the time it takes to receive results can be prolonged. Therefore, every parameter allowing for early diagnosis is vital. Therefore, in this study, the possibility of diagnosing COVID-19 early by a simple, inexpensive, easily accessible test, such as a CBC, has been examined.

Patients who came with history of fever and were admitted in the hospital, RTPCR was performed during the arrival and was found that 65% patients were covid positive and 35% patients were negative out of 100 patients examined.

In our study COVID 19 positive patients presented with leucopenia, neutropenia, thrombocytopenia, lymphocytosis, SII and NLR were reduced, PLR and CRP was Increased. PT and APTT were within normal limits. Lymphocyte count and PLR showed strong significance in this study. Comparison was made between current study and various other studies as described in Table 5 where concordance and discordance was seen between different studies.

Table 5: Comparison between current study and other study

Parameters	Current study	Eren Usul et al	Terpos et al	Guan et al	Liu et al	Huang et al.	Fan et al
Age	43.03±16.1	47.6± 16.9	-	-	-	-	-
Hemoglobin	Increased	Increased	-	-	-	-	-
Tlc	Decreased	Decreased	Reduced	Decreased	Decreased	Reduced	Decreased
Neutrophils	Decreased	Decreased	-	-	Increased	-	Decreased
Lymphocytes	Increased	Decreased	Reduced	Decreased	-	-	Decreased
Platelets	Decreased	Decreased	-	Decreased	Decreased	Decreased	Decreased
SII	Decreased	Decreased	-	-	-	-	-
NLR	Decreased	Increased	Increased	-	-	-	-
PLR	Increased	Decreased	-	-	-	-	-
PT	Normal	-	-	-	-	Increased	-
APTT	Normal	-	-	-	-	Decreased	-
CRP	Increased	-	-	Increased	Increased	Increased	-

**Fig. 1:** Receiver operating characteristic curve for significant markers in the prediction of corona virus positive patients

An obvious relationship, although not definitely proven, exists between a bacterial infection and neutrophilia, and a viral infection and lymphocytosis. Accordingly, NLR of peripheral blood has been used to distinguish between these types of infections. In few studies NLR is higher in those with fever due to bacterial infections than those with fever due to a viral etiology¹ however in our study NLR was reduced in COVID-19 positive cases. SII has been proposed as a prognostic indicator in the follow-up of sepsis patients.⁵ In this study, SII was found to be significantly low for COVID-19-positive patients, meaning that it can also be used while diagnosing COVID-19.

5. Conclusion

The definitive diagnosis of COVID-19 was made by RT-PCR analysis, but this is a time-consuming and less accessible test. In our study, low values of leukocytes, neutrophils, platelets and high values of hemoglobin, Lymphocytes were found with a CBC test which is easily

available are found to be valuable in terms of the initial diagnosis of COVID 19. In addition, low values of NLR and SII and high value of PLR and CRP are also indicative of COVID-19.

6. Source of Funding

None.


7. Conflict of Interest

None.

References

- Usul E, Şan İ, Bekgöz B, Şahin A. The role of hematological parameters in COVID-19 patients in the emergency room. *Biomark Med.* 2020;doi:10.2217/bmm-2020-0317.
- Fan BE. Hematologic parameters in patients with COVID19 infection. *Am J Hematol.* 2020;95(8):E215.
- Huang C. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497–506.
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA.* 2020;323(11):1061–9.
- Lagunas-Alvarado M, Mijangos-Huesca FJ, an Gonz' alez JOT. Systemic immune inflammatory index in sepsis. *Med Int Mex.* 2017;33(3):303–9.

Author biography

Sneha Aditi, Consultant Pathologist and Lab Head
 <https://orcid.org/0000-0003-2000-815X>

Roushni Kumari, Senior Resident

Abdul Aziz, Consultant Pathologist

Cite this article: Aditi S, Kumari R, Aziz A. The role of hematological biomarkers, coagulation profile and CRP (c-reactive protein) in COVID-19 patients in private health care sector. *Indian J Pathol Oncol* 2022;9(3):216-219.