

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pathology and Oncology

Journal homepage: www.ijpo.co.in

Editorial

Serrated lesions of colon

Bhanushree C S ¹*

¹Dept. of Pathology, Indira Gandhi Medical College and Research Institute, Kathirkamam, Puducherry, India



ARTICLE INFO

Article history:

Received 12-11-2023

Accepted 18-11-2023

Available online 11-12-2023

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

A broad spectrum of lesions ranging from benign to malignant found in colon. The lesions comprised of micro vesicular hyperplastic polyp, goblet cell rich hyperplastic polyp, sessile serrated lesion, sessile serrated lesion with dysplasia, traditional serrated adenoma and serrated carcinoma.¹ Serrated lesion simply means saw tooth morphology on surface of the polyp and crypt.² Serrated lesions are very difficult to diagnose on histology and its very important to diagnose such polyps as sessile serrated polyps can transform to carcinoma. They are more common on right side of colon and carry BRAF mutation.³ The progression of sessile serrated polyp to cancer follows serrated pathway. Histologically, marked serrations at the base of the crypt with distorted architecture will be seen.⁴ The traditional serrated adenoma also carries the risk of malignancy and have K-RAS mutation. Histologically, show villous or tubulovillous architecture having columnar cells with diffusely eosinophilic cytoplasm along with aberrant crypts.⁵ The three molecular features of serrated polyps are BRAF mutation, KRAS mutation and CpG island methylation. The sessile serrated polyp progress through BRAF serrated pathway and traditional Serrated polyps by KRAS serrated pathway accounting to 15-20% of colorectal cancer.⁶ The knowledge about serrated lesions aids in the better management of colorectal cancer.

References

1. Rosty C, Hewett DG, Brown IS, Leggett BA, Whitehall VL. Serrated polyps of the large intestine: current understanding of diagnosis, pathogenesis, and clinical management. *J Gastroenterol.* 2013;48(3):287–302.
2. Jaravaza DR, Rigby JM. Hyperplastic polyp or sessile serrated lesion? The contribution of serial sections to reclassification. *Diagn Pathol.* 2020;15(1):140. doi:10.1186/s13000-020-01057-0.
3. Rex DK. Serrated Polyps in the Colon. *Gastroenterol Hepatol (N Y).* 2014;10(10):671–4.
4. Monreal-Robles R, Jáquez-Quintana JO, Benavides-Salgado DE, González-González JA. Pólipos serrados del colon y el recto: una revisión concisa. *Rev de Gastroenterol Méx.* 2021;86(3):276–86.
5. Torlakovic EE, Gomez JD, Driman DK, Parfitt JR, Wang C, Benerjee T, et al. Sessile serrated adenoma (SSA) vs. traditional serrated adenoma (TSA). *Am J Surg Pathol.* 2008;32(1):21–9.
6. Kim SY, Kim TI. Serrated neoplasia pathway as an alternative route of colorectal cancer carcinogenesis. *Intest Res.* 2018;16(3):358–5.

Author biography

Bhanushree C S, Professor <https://orcid.org/0000-0003-4637-8837>

Cite this article: Bhanushree C S. Serrated lesions of colon. *Indian J Pathol Oncol* 2023;10(4):332-332.

* Corresponding author.

E-mail address: drbanushree15@hotmail.com (Bhanushree C S).