

Incidence of acute and chronic leukemias in rural area at tertiary care teaching hospital: a five years of study

Jaya Bhaskar Baviskar

Assistant Professor, Pravara Institute of Medical Sciences, Loni, Maharashtra

Email: drjayabaviskar@gmail.com

Abstract

Introduction: Leukemias are neoplastic proliferations of haematopoietic cells and form a major proportion of haematopoietic neoplasms that are diagnosed worldwide.

Aims: To chart out the incidence of four major types of leukemias presenting for the first time to Hospital located in rural region.

Methods: It was a prospective and retrospective study of 156 patients carried out in the department of Pathology located in rural area over a five years period between May 2006 and May 2011. Diagnosis was based on peripheral blood count, peripheral blood smear and bone marrow examination for morphology along with cytochemistry study whenever required.

Results: In this study, commonest leukemia was chronic myeloid leukemia (CML) followed by Acute lymphoid leukemia (ALL), Acute myeloid leukemia(AML) and then chronic lymphoid leukemia (CLL). Out of total 156 cases, 90 were male and 66 were females with Male: Female ratio is 1.36:1. Acute lymphoid leukemia was the most common type of leukaemia in the children and adolescents. Myeloid neoplasms were most common in adults.

Conclusion: CML was commonest type of leukemia followed by acute lymphoid leukemia. Male predominance was seen in all types of leukemia and age has significant effect on type of leukemia.

Keywords: Incidence, Leukemia, Rural India, Tertiary care teaching hospital

Introduction

The leukemias are defined as diseases in which abnormal proliferation of haematopoietic cells cause progressively increasing infiltration of bone marrow, although in certain forms the lymphatic tissues are particularly affected.¹ They form a significant percentage of haematological disorders and affects individuals of all age groups throughout the world, but the incidence of disease and the frequency of various morphological types and sub-types have been found to be differing in different countries.² Leukemias demonstrate extra-ordinary biological, morphological and clinical heterogeneity.³ Leukemias are usually divided into myeloid and lymphoid, acute and chronic. This simple classification based on lineage and on rate of disease progression when untreated. The purpose of classification of leukemias is to organise knowledge into manageable forms so that biological entities can be recognised. Correct classification permits optimal patient management.⁴

Leukemias are the 10th most common cancer in men and 12th most common in women and constitute 3% of the global cancer burden.⁵ Developing countries bear more than half of global cancer burden, because 75% of the world population lives in these countries.⁶ The incidence of Leukemia is highest in North America and Australia/ New-zealand and lowest in sub-saharan Africa.⁷ In India, lympho-haematopoietic malignancies constitute 9.5% of all cancers in men and 5.5% in women.⁸ As per available information from population based surveys, the incidence of leukemia in India varies from 0.8/1, 00,000 in Barshi (Rural area of Maharashtra) to 5/1, 00,000 in Delhi. These figures are comparably lower than rest of the world but under-

diagnosis and under-reporting cannot be ruled out.⁸ The cell type distribution of leukemias observed in India is different from that observed in developed world. Myeloid leukemias predominate in India while lymphoid leukemias dominate in western world mainly because of higher incidence of chronic lymphatic leukemia.² Despite being relative uncommon, leukemias have been studied more extensively because of easy accessibility of involved tissue.⁵ The incidence of CML was noted highest(45.3%) and that was lowest of CLL(5.7%) in Capital of India i.e. Delhi during a period of 1970-1979.⁹ Similar observations were noted in Chandigarh and other metrocities like Mumbai and Calcutta.^{10,11,12} There was exception for incidence of ALL(39.2%) which was highest observed in Kerala state during a period of 1980-1983.¹³

Hence this study was carried out in rural population to know that rural population differs markedly from rest of urban Indian population or not.

Materials and Methods

Present study was carried out in the department of pathology, during the period of May 2006- May 2011. A total of 156 patients with acute and chronic leukemia reported during the study period were included in the study. F.A.B. (French, American, British) classification was used for AML which subdivides into eight types M0, M1, M2, M3, M4, M5, M6, M7 and ALL into three subtypes as L1, L2, L3.^{14,15,16} These patients were evaluated especially regarding age, gender, risk factors, occupation and chief complaints.

The study from May 2006 to April 2009 was retrospective and data collected from records maintained in department of Pathology, Haematology

and Medical record section of same Hospital. All cases reporting to Hospital for the first time from May 2009 to May 2011 were prospective cases and their clinico-laboratory characteristics were recorded.

Diagnosis was made on findings of peripheral blood count (i.e. Haemoglobin, Total leucocyte count and Platelet count) which was done by automated cell counter; peripheral blood smear and bone marrow aspiration smear stained by Leishman stain. Leishman stain was poured on air dried unfixed smears for a period of 5-7 minutes, then added twice quantity of buffered distilled water for next 10 minutes.¹⁷ Cytochemistry studies (like myeloperoxidase (MPO), Periodic acid schiff stain (PAS), Sudan black B (SBB)

and Non-specific esterase) were done in all prospective cases to differentiate myeloid and lymphoid leukemia.¹⁷

These stains were prepared and used as per the guidelines endorsed in Practical Haematology Dacie and Lewis 10th edition, year 2006.¹⁷ Then all cases diagnosed and classified, as acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia and chronic lymphoid leukemias.

Results

Total 156 patients diagnosed as leukaemia were further subjected to analysis according to their different morphology and other relevant parameters.

Table 1: Gender wise distribution of different types of Leukaemia amongst study population

Gender	AML	ALL	CML	CLL	CMML	JMML	Total
Male	15 (9.61%)	23 (14.74%)	35 (22.43%)	15 (9.61%)	01 (0.64%)	01 (0.64%)	90 (57.70%)
Female	21 (13.46%)	18 (11.54%)	18 (11.54%)	09 (5.76%)	00	00	66 (42.30%)
Total	36 (23.07%)	41 (26.28%)	53 (33.97%)	24 (15.38%)	01 (0.64%)	01 (0.64%)	156 (100%)

(AML- Acute myeloid leukemia, ALL- Acute lymphoid leukemia, CML- Chronic myeloid leukemia, CLL- Chronic lymphoid leukemia, CMML- Chronic myelomonocytic leukemia, JMML- Juvenile myelomonocytic leukemia)

Table 1 shows that the commonest type of leukemia encountered was chronic myeloid leukemia (CML) which is accounted for 33.97% cases followed by acute lymphoid leukemia (ALL) (26.28%), acute myeloid leukemia (AML) (23.07%) and then chronic lymphoid leukemia (15.38%). It was also observed that, out of 156 cases, 90 were male and 66 were females and Male: Female ratio was 1.36:1 which indicates male predominance in all types of leukemia.

Table 2: Age and Sex distribution of leukemia amongst study population

Sex/Age	AML			ALL			CML			CLL		
	M	F	Total	M	F	Total	M	F	Total	M	F	Total
1-15	02	01	03	21	17	38	00	01	01	00	00	00
16-30	06	02	08	02	01	03	05	04	09	01	00	01
31-45	03	03	06	00	00	00	11	10	21	02	00	02
46-60	09	06	15	00	00	00	13	03	16	07	03	10
61-75	01	02	03	00	00	00	05	00	05	05	05	10
>75	00	01	01	00	00	00	01	00	01	00	01	01
Total	21	15	36	23	18	41	35	18	53	15	09	24
M:F Ratio	1.4:1			1.28:1			1.94:1			1.66:1		

It was observed from table 2 that age groups found to be most vulnerable for the various leukemias, which was as follows; AML was commonest in age group of 46-50 years, ALL were between 1-15 years of age group, and peak incidence found in the age group of 1-5 years followed by 6-10 years. CML were evenly distributed from the age group of 21-65 years; and peak incidence was found in between 31-45 years followed by 46-60 years, while CLL were more common in the age group of 61-75 year.

Discussion

In the year of 1857, Virchow probably was the first to classify leukemia. On the pathologic distribution of tumor he distinguished splenic and lymphatic forms of leukemia.¹⁸ The incidence of leukemia has increased considerably and this rise is noticeable because of improved statistics, better case findings with novel technologies which lead to better diagnosis and treatment methods. The incidence of leukemia found to be varying from different geographical areas according to their life styles, economic conditions, and poverty rate.¹⁹

The incidence and clinical presentation of many haematological cancers in India is relatively lower and differs from that seen in western world due to health awareness and availability of health care delivery system.^{20,21}

In present study, chronic myeloid leukemia (CML) was more common which was in concordance with the studies carried out by Vasavada *et al.*²², Kushawaha *et al.*²³, Chatterji *et al.*¹², Rani S *et al.*⁹, Shome *et al.*¹⁰ and Rathee *et al.*²⁴. It was also observed that studies reported by Menzes and Malik²⁵ and Verghese *et al.*¹³ vary from present study, which is might due to geographical variation and literate public. Chronic myeloid leukemia is unique in combining the need for an expensive long-term therapy (daily TKIs) that results in clinical elimination of the disease only with continued therapy in most instances making it more of chronic disease. Due to chronicity of CML incidence and prevalence remains higher amongst all leukemias. It is likely that patients with CML are now being diagnosed earlier for several reasons including greater awareness of the disease among clinicians and the wide availability of cheap, accurate leucocyte and platelet measurements not performed in the past.

In our study, male predominance was seen, which is similar with most studies conducted in India and other countries.^{24,26}

Present study found that acute lymphoid leukemia (ALL) was the second most common leukemia subtype which was in accordance with the studies reported by Modak H *et al.*²⁷, Chen *et al.*²⁸ and Shome *et al.*¹⁰. However our observations varies from the studies conducted by Vasavada *et al.*²² and Kushawaha *et al.*²³ which might be due to under diagnosis or under reporting. It is also observed that, maximum cases are seen in first and second decade of life which comparable with D'Costa GG *et al.*¹¹, Parkin DM *et al.*²⁹, Meighan *et al.*³⁰, Alrudainy *et al.*³¹ Male predominance for ALL was found in North India (Delhi), rural area of Maharashtra like Barshi, and South India (Chennai and Bangalore), which is in accordance with present study.³²

Acute myeloid leukemia (AML) was the third most common leukemia in the present study which was in accordance with study carried out by D'Costa GG *et al.*¹¹. Incidence of lower percentage was reported by Vasavada *et al.*²² and Menzes and Malik²⁵ which was vary from the present study. Large percentage of cases found in adults was comparable with most of studies in India, eastern and western countries. In this study male predominance was seen, which was similar with most studies mentioned.^{27,33-36}

Though, CLL is the most common form of leukemia affecting adults in western countries. Incidence of CLL is lower in India (1.95-8.8%)^{9-13,23,37}. In present study, chronic lymphoid leukemia (CLL) has lowest percentage of total leukemia cases. As the overall incidence of CLL is lower in India, bit higher

incidence was observed in present study. It was also observed that most of the cases of CLL (70.83%) were seen in patients more than 55 years of age and it is comparable with the study performed in AIIMS Hospital (68.42%).³⁸

In present study, Juvenile Myelomonocytic leukemia (JMML) & Chronic myelomonocytic leukemia (CMML) types of leukaemia found to be only a single case respectively as the overall incidence is very rare for these leukaemias, which might be related to our geographical area.

Conclusion

Present study concluded that chronic leukemia was more common than acute leukemia. The most common type of leukemia was CML followed by ALL, AML and CLL. Male predominance was seen in all types of leukemia. Age was important factor, as specific age groups were observed more common in some types of leukemias. In children, ALL was more common, while AML was more commonly seen in adults. CML and CLL were observed only in adults. CLL was more commonly observed in elderly patients. Present study concluded that the incidence of different types of leukaemia from rural populations doesn't differ markedly from rest of the urban Indian populations.

References

1. Robbins and Cotran: Pathologic Basis of Diseases, 7th Edition. Elsevier Publications, 1999. Pg. 232-37.
2. Shashi Prakash, Rama Murthi, Rama Gopalan Aurora AI, Leukemias at Pondicherry, Indian J cancer 1981;18:1-6.
3. Causar J B, The haematopoietic- lymphoid Neoplasms chapter 80. In Wintrob's clinical haematology. Lee G.R. *et al*, Baltimore: Williams and Wilkins 1998.
4. J.A. Whittaker and J.A. holmes Leukemia and related disorders 3rd edition, Blackwell science publications, page 31.
5. Parkin OM, Pisani P, Ferlay J. Estimate of the world wide frequency of eighteen major cancer in 1995, Int J cancer 1993;54:504-606.
6. WHO 1999. Health situation in the South East Asia Region 1994-1997, Regional Office for SEAR, New Delhi, 1999.
7. Parkin DM, Whelam SL, Ferlay J, Raymond L, Young J, eds., Cancer incidence in five continents, Vol VII IARC Scientific publications number 143 Lyon 1997, International agency for research on cancer pp 804-913.
8. National cancer Registry programme, consolidated report of population based cancer registries 1990-96. Incidence and distribution of cancer Indian council of Medical Research, New Delhi, Aug 2001.
9. Rani S, Beohar PC, Mohanty TK, Mathur MD, Leukemic Pattern in Delhi – A Ten year study of 490 cases, Indian J of Cancer, 1982;19:81-86.
10. Shome DK, Ghosh K, Mohanty D, Das K, Leukaemia in North West India, Acta Haematol, 1985;73:244.
11. Dicosta GG, Siddique HM, Pradhan RM, Gupte SS, Pattern of leukemias: a ten year incidence study of 242 cases, J Postgraduate Med 1989;35:191-95.
12. Chatterji JB, Ghose S, Ray RN, Incidence of Leukemia. An analysis of 544 cases studied in Calcutta, J Assoc Phy Ind, 1962;10:673-76.

13. Varghese PR, Elayidom NB, Joseph CD, Kumar S. Epidemiological observations on leukaemia in Kerala (A study of 1016 cases over three years), *Ind J Haematol*, 1984;2:15-17.
14. Firkin F., Chesterman C., Penington D., Rush B. — DeGruchy's Clinical Haematology in Medical Practice. 5th ed. Osney Mead Oxford OX2 0EL: Blackwell Science Ltd. P 236-277, 1989.
15. Cousar J.B. — Hematopoietic-Lymphoid neoplasm: Principle of Diagnosis. In: Greer JP, Foerster J, Rodgers GM, Paraskevas F, Glader B editors. *Wintrobe Clinical Haematology*, 11th Edn. Lippincott Williams & Wilkins: Philadelphia. p. 1913-1917, 2004.
16. Bain B.J., Bates I. — Approach to the diagnosis and classification of blood diseases. In: Lewis SM, Bain BJ, Bates I editors. *Practical Haematology* 10th ed. Churchill Livingstone: Elsevier Ltd. p.609-624, 2006.
17. *Practical Haematology* Dacie and Lewis, 10th edition, Year 2006, page 65, 318-326.
18. Gralnick H R, Galton DAG, Catavasky D, Sultan C, Bennett J M, Classification of Acute leukemia, *Annals of Internal Medicine* 1977;87:740-53.
19. Gunz F.W., The epidemiology and genetics of chronic leukemias, *Clinical Haematology*, 1977;6:3-20.
20. Kumar L, Kumari M, Kumar S, Kochupillai V, Singh R, Clinical and laboratory features at diagnosis in 437 patients with chronic myelogenous leukemia: An experience at tertiary care centre, In: Kumar L(ed). *Progress in hematologic oncology*, New York: The advanced research foundation 2003;83-98.
21. Kulshreshta R, Sah SP- Pattern of occurrence of leukemia at a teaching Hospital in Eastern region of Nepal- A posted the SEER web site, 2010 six year study *J Nepal Med Assoc* 2009;48(173):35-40.
22. Vasavada PS, Akbar Ali and DP Mukherjee Clinical studies in leukemia- A review of 100 cases in symposium in leukemia. *Leukemia journal of associations Phy India*, 1962;9:357.
23. Kushwaha MRS, Bagchi M, Mehrotra RML, Leukemia at Lucknow- a study of 200 cases, *Ind J Cancer* 1978;15:28-34.
24. Radha Rathee, Minakshi Vashist, Ashok Kumar, Sunita Singh, Incidence of acute and chronic forms of leukemia in Haryana; *International Journal of Pharmacy and Pharmaceutical Sciences*; 2014;6(2):42-45.
25. Menzes S and GB Malik, A study of leukemias in Goa- *Indian J of Path and Micro*, 1977;20:83-90.
26. Wing Y. Au Priscilla B. Caguioa Charles Chuah Szu Chun Hsu Saengsuree Jootar Dong- Wook Kim II- Young Kweon William M. O'Neil Tapan K. Saikia Jianxiang Wang: Chronic myeloid leukemia in Asia; *Int J Haematol* 2009;89:14-23.
27. Modak H., Kulkarni S.S., Kadakol G.S., Hiremathb S.V., Patil B.R., Hallikeri U. et al. Prevalence and Risk of Leukemia in the Multi-ethnic Population of North Karnataka. *Asian Pacific J Cancer Prev* 2011;12:671-75.
28. Chen B., Huang Z., Zhang X., Ou-Yang J., Li J., Zhai Y. et al. An epidemiological investigation of leukemia incidence between 2003 and 2007 in Nanjing, China. *J Hematol Oncol* 2010;3:21.
29. Parkin D.M., Stiller C.A., Draper G.J., Bieber C.A. The international incidence of Childhood cancer. *Int J Cancer* 1988;42(4):511-20.
30. Meighan S.S. Leukemia in children. Incidence, clinical manifestation, and survival in an unselected series, *J Amer Med Assoc*, 1964;190:578-582.
31. Alrudainy L.A., Salih H.M., Aldorky M.K. Incidence and Pattern of Childhood Leukemia in Basrah, Iraq during 2003-2007 *IJBC* 2009;5:11-17.
32. Arora RS, Eden TOB, Kapoor G: Epidemiology of childhood cancer in India; *Indian journal of cancer* 2009;46(4):27-32.
33. Firkin F., Chesterman C., Penington D., Rush B. DeGruchy's Clinical Haematology in Medical Practice, 5th ed., Osney Mead Oxford OX2 0EL: Blackwell Science Ltd. 1989;236-277.
34. Ferlay J., Shin H.R., Bray F., Forman D., Mathers C., Parkin D.M. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008, (2010) UICC. *Int. J. Cancer* 2010;127:2893–2917.
35. Gosh S., Shinde S.C., Kumaran G.S., Sapre R.S., Dhond S.R., Badrinath Y. et al. Hematologic and Immunophenotypic Profile of Acute Myeloid Leukemia: An Experience of Tata Memorial Hospital. *Indian J Cancer* 2003;40(2):71-76.
36. Eivazi-Ziaei J. Myeloperoxidase Index and Subtypes of Acute Myeloid Leukemia. *J Pak Med Assoc* 2009;59(6):406-07.
37. Advani SH, Jussawala DJ, Nagaraj RD, Gangadharan P, Shetty PA, A study of 1126 leukemia cases- Epidemiologic and end result analysis, *The Indian J of Cancer* 1979;16:8–17.
38. Neeraja Agrawal, Rahul Naithani, M. Mahapatra, Inusha Panigrahi, Rajat Kumar, H.P. Pati, Renu Saxena, V.P. Chaudhary, Chronic lymphocytic leukemia in India- A clinicopathological profile: *Hematology* 2007;12(3):229-33.